

***Self-consistent Green's function in
Finite Nuclei and related things...***

-

Lecture III

***Algebraic Diagrammatic Construction method
(ADC) for calculating the self- energy***



Current Status of low-energy nuclear physics

Composite system of interacting fermions

Binding and limits of stability

Coexistence of individual and collective behaviors


Self-organization and emerging phenomena

EOS of neutron star matter

Experimental

programs

RIKEN, FAIR, FRIB



II) Nuclear correlations

Fully known for stable isotopes

[C. Barbieri and W. H. Dickhoff, Prog. Part. Nucl. Phys 52, 377 (2004)]

Neutron-rich nuclei; Shell evolution (far from stability)

I) Understanding the nuclear force

QCD-derived; 3-nucleon forces (3NFs)

First principle (ab-initio) predictions

III) Interdisciplinary character

Astrophysics

Tests of the standard model

Other fermionic systems:

ultracold gasses; molecules;

Nuclear forces in exotic nuclei

Nucleon interactions are very complex and difficult to handle

Change of regime from stable to dripline isotopes !


Symmetric matter:
 $N \approx Z$





Neutron-rich matter ($N \gg Z$):

- Neutron star matter EoS
- Symmetry energy
- New shell closures


Tensor force (p-n)



Three-nucleon Force (3NF)



Driplines of nitrogen and fluorine isotopes




[A. Cipollone, CB, P. Navrátil, Phys. Rev. Lett. **111**, 062501 (2013)]

Green's functions in many-body theory

One-body Green's function (or propagator) describes the motion of quasi-particles and holes:

$$g_{\alpha\beta}(E) = \sum_n \frac{\langle \Psi_0^A | c_\alpha | \Psi_n^{A+1} \rangle \langle \Psi_n^{A+1} | c_\beta^\dagger | \Psi_0^A \rangle}{E - (E_n^{A+1} - E_0^A) + i\eta} + \sum_k \frac{\langle \Psi_0^A | c_\beta^\dagger | \Psi_k^{A-1} \rangle \langle \Psi_k^{A-1} | c_\alpha | \Psi_0^A \rangle}{E - (E_0^A - E_k^{A-1}) - i\eta}$$

...this contains all the structure information probed by nucleon transfer (spectral function):




Example of spectral function ^{56}Ni

One-body Green's function (or propagator) describes the motion of quasi-particles and holes:

$$g_{\alpha\beta}(E) = \sum_n \frac{\langle \Psi_0^A | c_\alpha | \Psi_n^{A+1} \rangle \langle \Psi_n^{A+1} | c_\beta^\dagger | \Psi_0^A \rangle}{E - (E_n^{A+1} - E_0^A) + i\eta} + \sum_k \frac{\langle \Psi_0^A | c_\beta^\dagger | \Psi_k^{A-1} \rangle \langle \Psi_k^{A-1} | c_\alpha | \Psi_0^A \rangle}{E - (E_0^A - E_k^{A-1}) - i\eta}$$

...this contains all the structure information probed by nucleon transfer (spectral function):

$$S_{ab}^h(\omega) = \frac{1}{\pi} \text{Im} g_{ab}(\omega)$$



Why want to look at spectral function?


- Strong insight in to the structure (can be done the “ab-initio” way but it also gives qualitative understanding)
- Koltun SR links E_0^A and S^h in a “deep” manner
- Describes hole states (structure) and particle region (scattering) naturally
- Response to excitations, and particle addition/removal
- Useful to investigate changes shell structure

Dyson equation

Dyson equation:

$$g_{\alpha\beta}(t - t') = g_{\alpha\beta}^{(0)}(t - t') + g_{\alpha\gamma}^{(0)}(t - t_\gamma) \Sigma_{\gamma\delta}^*(t_\gamma, t_\delta) g_{\delta\beta}(t_\gamma - t')$$

Diagrammatically:



Approaches to compute the irreducible self-energy:

- Use PT → Feynman diagram expansion
- Equation of Motion method
 - Leads to important concepts:
 - self consistency
 - all-order summations
 - conservation theorems
- Algebraic diagrammatic constructions ADC(3)
 - typically the working approach for most finite systems

Adiabatic theorem and perturbations

Assume that the Hamiltonian splits in two parts, one component (H_0) can be solved exactly but not the full Hamiltonian:

$$H = H_0 + H_1$$

If the second part (H_1) is small, we can treat it as a small correction \rightarrow *perturbation theory*.

The complete propagator requires the Heisenberg evolution for the full H :

$$g_{\alpha\beta}(t, t') = -\frac{i}{\hbar} \langle \Psi_0^N | T[c_\alpha(t) c_\beta^\dagger(t')] | \Psi_0^N \rangle, \quad c_\alpha(t) = e^{iHt/\hbar} c_\alpha e^{-iHt/\hbar}$$

but we can handle only H_0 . Thus, evolve operators according to an (H_0) compensate for the missing part (H_1) evolving the wave function \rightarrow This is the *Interaction (or Dirac) picture*.

Feynman diagram rules

Graphic conventions:

$$i\hbar g_{\alpha\beta}(t-t') = \begin{array}{c} \alpha \\ \parallel \\ \blacktriangleright \\ \parallel \\ \beta \end{array} = i\hbar g_{\alpha\beta}(\omega)$$

$$\frac{-i}{\hbar} u_{\alpha\beta}, \quad \frac{-i}{\hbar} t_{\alpha\beta} = \begin{array}{c} \alpha \\ \bullet \\ \text{---} \\ \beta \end{array} \quad \times$$

$$i\hbar g_{\alpha\beta}^{(0)}(t-t') = \begin{array}{c} \alpha \\ \diagup \\ \blacktriangleright \\ \diagdown \\ \beta \end{array} = i\hbar g_{\alpha\beta}^{(0)}(\omega)$$

$$\frac{-i}{\hbar} v_{\alpha\beta,\gamma\delta} = \begin{array}{c} \alpha \\ \bullet \\ \text{---} \\ \gamma \end{array} \quad \begin{array}{c} \beta \\ \bullet \\ \text{---} \\ \delta \end{array}$$

$$i\hbar g_{\alpha\beta\gamma\dots;\delta\dots}(t_\alpha, t_\beta, \dots; t_\delta) = \begin{array}{c} \alpha \quad \beta \quad \gamma \quad \dots \\ \blacktriangleright \quad \blacktriangleright \quad \blacktriangleright \quad \dots \\ \text{---} \\ g^{4\text{-pt}} \\ \text{---} \\ \blacktriangleright \quad \dots \\ \delta \end{array} = i\hbar g_{\alpha\beta\gamma\dots;\delta\dots}(\omega_\alpha, \omega_\beta, \dots; \omega_\delta)$$

Feynman diagram rules

Rules in *time* representation

1. Write all **connected** and **topologically equivalent** diagrams—and only those.
2. Each single line w/ an arrow, contributes $\frac{1}{i} \int_{\alpha}^{\beta} dt$ running from β to α
3. Each closed circle contributes a density matrix $\rho(\alpha, \alpha)$ (no i factor!)
4. Each two-body interaction line contributes $\frac{1}{i} \int dt V(t)$
5. Each external field line contributes $\int dt \phi(t)$
6. Add an extra **-1** factor for each closed circuit (the density matrix loops **excluded**)
7. Sum (integrate) over all coordinate and integrate over all internal times
8. IF ψ are antisymmetrized matrix elements, and extra factor $\frac{1}{2}$ is required for each pair of equivalent lines, starting from the **common** interaction and ending on **common** interaction (not necessarily the same).
9. Add final factor i^n is to get $G(t-t')$.

Feynman diagram rules

One can transform any propagator in frequency space.
This is done by:

$$g_{\alpha\beta\dots;\mu\nu\dots}(\omega_\alpha, \omega_\beta, \dots) = \int dt_\alpha \int dt_\beta \cdots \int dt_\mu \int dt_\nu \cdots \\ \times e^{i\omega_\alpha t_\alpha} e^{i\omega_\beta t_\beta} \cdots g_{\alpha\beta\dots;\mu\nu\dots}(t_\alpha, t_\beta, \dots; t_\mu, t_\nu, \dots) e^{-i\omega_\mu t_\mu} e^{-i\omega_\nu t_\nu} \cdots$$

Note that:

$$g_{\alpha\beta}(\omega, \omega') = \delta(\omega - \omega') g_{\alpha\beta}(\omega)$$

usual transformation for
the 2-time propagators.

For the interactions: after Fourier transformation the delta terms in $t_{\alpha\beta}\delta(t_\alpha - t_\beta)$ and $v_{\alpha\beta,\gamma\delta}\delta(t_\alpha - t_\beta)\delta(t_\gamma - t_\delta)\delta(t_\alpha - t_\gamma)$ give the conservation of incoming and outgoing energy.

Feynman diagram rules


Rules in *frequency* representation

1. Write all **connected** and **topologically equivalent** diagrams—and only those.
2. At every propagator line one must associate an energy going in the direction of the arrow (energy must be conserved at each vertex)
3. Each single line w/ an arrow, contributes $i\hbar g_{\alpha\beta}^{(0)}(\omega)$ running from β to α (ω gets a - sign if it goes against the arrow)
4. Each closed circle contributes a density matrix $\rho_{\alpha\beta}$ (no $i\hbar$ factor!)
5. Each two-body interaction line contributes $-\frac{i}{\hbar} v_{\alpha\beta,\gamma\delta}$
6. Each external field line contributes $-\frac{i}{\hbar} u_{\alpha\beta}$
7. An extra **-1** for each closed circuit (density matrix loops **excluded**)
8. Sum (integrate) over all coordinate and integrate over all independent frequencies (with a $1/2\pi$ factor for each integration)
9. IF $v_{\alpha\beta,\gamma\delta}$ are antisymmetrized matrix elements, and extra factor $\frac{1}{2}$ is required for each pair of equivalent lines.
10. Add final factor $-\frac{i}{\hbar}$ is to get $G(t-t')$.

Example of using Feynman diagram rules

Calculating the *second order self-energy*:

$$\Sigma_{\alpha\beta}^{2nd}(\omega) = i\hbar (i\hbar)^3 \left(\frac{-i}{\hbar}\right)^2 \frac{(-1)^2}{2} v_{\alpha\gamma,\kappa\zeta} v_{\mu\nu,\beta\lambda} \int \frac{d\omega_1}{2\pi} \int \frac{d\omega_2}{2\pi} g_{\kappa\mu}(\omega - \omega_1) g_{\zeta\nu}(\omega_1 + \omega_2) g_{\lambda\gamma}(\omega_2)$$



Repeated greek indices are implicitly summed


Example of using Feynman diagram rules

Calculating the *second order self-energy*:

$$x_i \equiv \hbar\omega_i$$

$$\begin{aligned} \Sigma_{\alpha\beta}^{2nd}(\omega) &= i\hbar (i\hbar)^3 \left(\frac{-i}{\hbar}\right)^2 \frac{(-1)^2}{2} v_{\alpha\gamma,\kappa\zeta} v_{\mu\nu,\beta\lambda} \int \frac{d\omega_1}{2\pi} \int \frac{d\omega_2}{2\pi} g_{\kappa\mu}(\omega - \omega_1) g_{\zeta\nu}(\omega_1 + \omega_2) g_{\lambda\gamma}(\omega_2) \\ &= -\frac{1}{2} v_{\alpha\gamma,\kappa\zeta} v_{\mu\nu,\beta\lambda} \int \frac{dx_1}{2\pi i} \int \frac{dx_2}{2\pi i} g_{\kappa\mu}(\omega - \omega_1) g_{\zeta\nu}(\omega_1 + \omega_2) g_{\lambda\gamma}(\omega_2) \\ &= -\frac{1}{2} v_{\alpha\gamma,\kappa\zeta} v_{\mu\nu,\beta\lambda} \int \frac{dx_1}{2\pi i} g_{\kappa\mu}(\omega - \omega_1) \int \frac{dx_2}{2\pi i} \\ &\quad \times \delta_{\zeta\nu} \left\{ \frac{\delta_{\zeta \notin F}}{x_1 + x_2 - \varepsilon_{\zeta}^+ + i\eta} + \frac{\delta_{\zeta \in F}}{x_1 + x_2 - \varepsilon_{\zeta}^- - i\eta} \right\} \left\{ \frac{\delta_{\gamma \notin F}}{x_2 - \varepsilon_{\gamma}^+ + i\eta} + \frac{\delta_{\gamma \in F}}{x_2 - \varepsilon_{\gamma}^- - i\eta} \right\} \end{aligned}$$

Using the Cauchy theorem, only term with at least one pole on each side of the real axis contribute:




Example of using Feynman diagram rules

Calculating the *second order self-energy*:

$$x_i \equiv \hbar\omega_i$$

$$\Sigma_{\alpha\beta}^{2nd}(\omega) = -\frac{1}{2}v_{\alpha\gamma,\kappa\zeta}v_{\mu\nu,\beta\lambda} \int \frac{dx_1}{2\pi i} g_{\kappa\mu}(\omega - \omega_1)$$

$$\delta_{\zeta\nu}\delta_{\lambda\gamma} \left\{ \frac{\delta_{\zeta \notin F} \delta_{\gamma \in F}}{x_1 - (\varepsilon_{\zeta}^+ - \varepsilon_{\gamma}^-) + i\eta} - \frac{\delta_{\zeta \in F} \delta_{\gamma \notin F}}{x_1 + (\varepsilon_{\gamma}^+ - \varepsilon_{\zeta}^-) - i\eta} \right\}$$



$$\Pi_{\zeta\gamma,\nu\lambda}^{(0)}(\omega_1)$$

Example of using Feynman diagram rules

Calculating the *second order self-energy*:


$$x_i \equiv \hbar\omega_i$$

$$\begin{aligned} \Sigma_{\alpha\beta}^{2nd}(\omega) &= -\frac{1}{2}v_{\alpha\gamma,\kappa\zeta}v_{\mu\nu,\beta\lambda} \int \frac{dx_1}{2\pi i} g_{\kappa\mu}(\omega - \omega_1) \\ &\quad \delta_{\zeta\nu}\delta_{\lambda\gamma} \left\{ \frac{\delta_{\zeta \notin F} \delta_{\gamma \in F}}{x_1 - (\varepsilon_{\zeta}^+ - \varepsilon_{\gamma}^-) + i\eta} - \frac{\delta_{\zeta \in F} \delta_{\gamma \notin F}}{x_1 + (\varepsilon_{\gamma}^+ - \varepsilon_{\zeta}^-) - i\eta} \right\} \\ &= -\frac{1}{2}\delta_{\kappa\mu}\delta_{\zeta\nu}\delta_{\lambda\gamma}v_{\alpha\gamma,\kappa\zeta}v_{\mu\nu,\beta\lambda} \int \frac{dx_1}{2\pi i} \left\{ \frac{\delta_{\kappa \notin F}}{\hbar\omega - \underset{x_1 \uparrow}{x_1} - \varepsilon_{\kappa}^+ + i\eta} + \frac{\delta_{\kappa \in F}}{\hbar\omega - \underset{x_1 \downarrow}{x_1} - \varepsilon_{\kappa}^- - i\eta} \right\} \\ &\quad \times \left\{ \frac{\delta_{\zeta \notin F} \delta_{\gamma \in F}}{\underset{x_1 \downarrow}{x_1} - (\varepsilon_{\zeta}^+ - \varepsilon_{\gamma}^-) + i\eta} - \frac{\delta_{\zeta \in F} \delta_{\gamma \notin F}}{\underset{x_1 \uparrow}{x_1} + (\varepsilon_{\gamma}^+ - \varepsilon_{\zeta}^-) - i\eta} \right\} \\ \Sigma_{\alpha\beta}^{2nd}(\omega) &= \frac{1}{2}v_{\alpha\lambda,\mu\nu} \left\{ \frac{\delta_{\mu \notin F} \delta_{\nu \notin F} \delta_{\lambda \in F}}{\hbar\omega - (\varepsilon_{\mu}^+ + \varepsilon_{\nu}^+ - \varepsilon_{\lambda}^-) + i\eta} + \frac{\delta_{\mu \in F} \delta_{\nu \in F} \delta_{\lambda \notin F}}{\hbar\omega - (\varepsilon_{\mu}^- + \varepsilon_{\nu}^- - \varepsilon_{\lambda}^+) - i\eta} \right\} v_{\mu\nu,\beta\lambda} \end{aligned}$$

Example of using Feynman diagram rules

Calculating the *second order self-energy*:

$$\Sigma_{\alpha\beta}^{2nd}(\omega) = i\hbar (i\hbar)^3 \left(\frac{-i}{\hbar}\right)^2 \frac{(-1)^2}{2} v_{\alpha\gamma,\kappa\zeta} v_{\mu\nu,\beta\lambda} \int \frac{d\omega_1}{2\pi} \int \frac{d\omega_2}{2\pi} g_{\kappa\mu}(\omega - \omega_1) g_{\zeta\nu}(\omega_1 + \omega_2) g_{\lambda\gamma}(\omega_2)$$



$$\Sigma_{\alpha\beta}^{2nd}(\omega) = \frac{1}{2} v_{\alpha\lambda,\mu\nu} \left\{ \frac{\delta_{\mu \notin F} \delta_{\nu \notin F} \delta_{\lambda \in F}}{\hbar\omega - (\varepsilon_{\mu}^+ + \varepsilon_{\nu}^+ - \varepsilon_{\lambda}^-) + i\eta} + \frac{\delta_{\mu \in F} \delta_{\nu \in F} \delta_{\lambda \notin F}}{\hbar\omega - (\varepsilon_{\mu}^- + \varepsilon_{\nu}^- - \varepsilon_{\lambda}^+) - i\eta} \right\} v_{\mu\nu,\beta\lambda}$$



Algebraic Diagrammatic Construction method at order n - $ADC(n)$

See J. Schirmer and collaborators.:

Phys. Rev. A**26**, 2395 (1982)

Phys. Rev. A**28**, 1237 (1983)

Working eqs. for ADC(2) / ext-ADC(2) / ADC(3)

We consider a generic *reference propagator* that is used to expand the self-energy:

$$g_{\alpha\beta}^{(ref)}(\omega) = \sum_n \frac{(\mathcal{X}_\alpha^n)^* \mathcal{X}_\beta^n}{\omega - \varepsilon_n^+ + i\eta} + \sum_k \frac{\mathcal{Y}_\alpha^k (\mathcal{Y}_\beta^k)^*}{\omega - \varepsilon_k^- - i\eta}$$

with

$$\left\{ \begin{array}{l} \mathcal{X}_\alpha^n \equiv \langle \Psi_n^{A+1} | a_\alpha^\dagger | \Psi_0^A \rangle \\ \varepsilon_n^+ \equiv E_n^{A+1} - E_0^A \\ E_n^{A+1} | \Psi_n^{A+1} \rangle = H^{(ref)} | \Psi_n^{A+1} \rangle \end{array} \right. \quad \left\{ \begin{array}{l} \mathcal{Y}_\alpha^k \equiv \langle \Psi_k^{A-1} | a_\alpha | \Psi_0^A \rangle \\ \varepsilon_k^- \equiv E_0^A - E_k^{A-1} \\ E_k^{A-1} | \Psi_k^{A-1} \rangle = H^{(ref)} | \Psi_k^{A-1} \rangle \end{array} \right.$$

In general, this could be an unperturbed propagator (for which $H^{(ref)} = H_0$, $\mathcal{X}_\alpha^n = \delta_{n,\alpha} \delta_{n \in F}$, etc...), an *Hartree-Fock propagator* or even fully dressed propagator.

Working eqs. for ADC(2) / ext-ADC(2) / ADC(3)

The most general form of the irreducible self-energy is:

$$\Sigma_{\alpha,\beta}^*(\omega) = \Sigma_{\alpha,\beta}^\infty + \sum_{ij} \mathbf{M}_{\alpha i}^\dagger \left[\frac{1}{\omega - (\mathbf{E}^{fw} + \mathbf{C}) + i\eta} \right]_{ij} \mathbf{M}_{j\beta} \\ + \sum_{rp} \mathbf{N}_{\alpha r}^\dagger \left[\frac{1}{\omega - (\mathbf{E}^{bk} + \mathbf{D}) - i\eta} \right]_{rp} \mathbf{N}_{p\beta}$$

where:

$i, j \longrightarrow$ label $2p1h, 3p2h, 4p3h, \dots$ excitations

$r, p \longrightarrow$ label $2h1p, 3h2p, \dots$ excitations

Working eqs. for ADC(2) / ext-ADC(2) / ADC(3)

The Dyson eq. is the solved by diagonalizing

$$\varepsilon^\pm \begin{pmatrix} \vec{Z}^\pm \\ \vec{W} \\ \vec{U} \end{pmatrix} = \begin{pmatrix} H_0 + \Sigma^\infty & \mathbf{M}^\dagger & \mathbf{N}^\dagger \\ \mathbf{M} & \text{diag}(\mathbf{E}^{fw}) + \mathbf{C} & \\ \mathbf{N} & & \text{diag}(\mathbf{E}^{bk}) + \mathbf{D} \end{pmatrix} \begin{pmatrix} \vec{Z}^\pm \\ \vec{W} \\ \vec{U} \end{pmatrix}$$

with the normalization condition

$$(\vec{Z}^\pm)^\dagger \vec{Z}^\pm + \vec{W}^\dagger \vec{W} + \vec{U}^\dagger \vec{U} = 1$$

One then identifies: $(\vec{Z}^{+n})_\alpha \rightarrow \mathcal{X}_\alpha^n$ that yield the new propagator and spectral function
 $(\vec{Z}^{-k})_\alpha \rightarrow \mathcal{Y}_\alpha^k$

Working eqs. for ADC(2)

The dressed 1st and 2nd order diagrams are:

$$\Sigma_{\alpha\beta}^{\infty} = \text{---} \times \quad \text{---} \circlearrowright = -U_{\alpha\beta} + \Sigma_{\alpha\beta}^{cHF}$$

$$\Sigma_{\alpha\beta}^{cHF} = \int_{C_{\uparrow}} \frac{d\omega}{2\pi i} v_{\alpha\gamma,\beta\delta} g_{\delta\gamma}^{(ref)}(\omega) = \sum_k v_{\alpha\gamma,\beta\delta} \mathcal{Y}_{\delta}^k (\mathcal{Y}_{\gamma}^k)^*$$

and

$$\Sigma_{\alpha\beta}^{(2)}(\omega) = \text{---} \circlearrowright + \text{---} \circlearrowright + \frac{1}{2} \sum_{n_1, n_2, k} \frac{v_{\alpha\lambda, \mu\nu} (\mathcal{X}_{\mu}^{n_1} \mathcal{X}_{\nu}^{n_2} \mathcal{Y}_{\lambda}^k)^* \mathcal{X}_{\mu'}^{n_1} \mathcal{X}_{\nu'}^{n_2} \mathcal{Y}_{\lambda'}^k v_{\mu'\nu', \beta\lambda'}}{\omega - (\varepsilon_{n_1}^+ + \varepsilon_{n_2}^+ - \varepsilon_k^-) + i\eta}$$

$$+ \frac{1}{2} \sum_{k_1, k_2, n} \frac{v_{\alpha\lambda, \mu\nu} \mathcal{Y}_{\mu}^{k_1} \mathcal{Y}_{\nu}^{k_2} \mathcal{X}_{\lambda}^n (\mathcal{Y}_{\mu'}^{k_1} \mathcal{Y}_{\nu'}^{k_2} \mathcal{X}_{\lambda'}^n)^* v_{\mu'\nu', \beta\lambda'}}{\omega - (\varepsilon_{k_1}^- + \varepsilon_{k_2}^- - \varepsilon_n^+) + i\eta}$$

Goldstone diagrams

Repeated greek indices are implicitly summed

Working eqs. for **ADC(2)**

From the previous diagrams, one extracts the matrix elements that define ADC(2):

$$\begin{aligned}(\mathbf{H}_0 + \Sigma^\infty)_{\alpha\beta} &= (\mathbf{T} + \mathbf{U})_{\alpha\beta} + (-\mathbf{U} + \Sigma^{cHF})_{\alpha\beta} \\ &= t_{\alpha\beta} + \sum_k v_{\alpha\gamma,\beta\delta} \mathcal{Y}_\delta^k (\mathcal{Y}_\gamma^k)^*\end{aligned}$$

$$\mathbf{M}_{(n_1,n_2,k),\alpha} = \frac{1}{\sqrt{2}} \mathcal{X}_\mu^{n_1} \mathcal{X}_\nu^{n_2} \mathcal{Y}_\lambda^k v_{\mu\nu,\alpha\lambda} \quad \mathbf{N}_{(k_1,k_2,n),\alpha} = \frac{1}{\sqrt{2}} (\mathcal{Y}_\mu^{k_1} \mathcal{Y}_\nu^{k_2} \mathcal{X}_\lambda^n)^* v_{\mu\nu,\alpha\lambda}$$

$$\mathbf{E}_{n_1,n_2,k}^{fw} = \varepsilon_{n_1}^+ + \varepsilon_{n_2}^+ - \varepsilon_k^-$$

$$\mathbf{E}_{k_1,k_2,n}^{bk} = \varepsilon_{k_1}^- + \varepsilon_{k_2}^- - \varepsilon_n^-$$

$$\mathbf{C} = 0$$


$$\mathbf{D} = 0$$

Any repeated indices are implicitly summed

Note that the auxiliary potential \mathbf{U} (that defines the unperturbed propagator) cancels out from the Dyson equation!

Working eqs. for *ext-ADC(2)*

Extend the ADC(2) by inserting pp-, hh-, and ph- summations (ladders and rings):



this leads to contributions of the form:

$$\begin{aligned}
 \rightarrow & V \frac{1}{\omega - E_{2p1h}} V + V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V \\
 & + V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V \\
 & + V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V \frac{1}{\omega - E_{2p1h}} V + \dots
 \end{aligned}$$

Working eqs. for *ext-ADC(2)*

Expand the self-energy in the inter-particle interaction. Both the \mathbf{M} , \mathbf{N} matrices have leading contributions at first order in V :

$$\mathbf{M} = \mathbf{M}^1(v^1) + \mathbf{M}^2(v^2) + \mathbf{M}^3(v^3) + \dots$$

$$\mathbf{N} = \mathbf{N}^1(v^1) + \mathbf{N}^2(v^2) + \mathbf{N}^3(v^3) + \dots$$

While \mathbf{C} and \mathbf{D} are only at 1st order in V . This leads to contributions of the form:

$$\begin{aligned} \mathbf{M}^\dagger \frac{1}{\omega - (E + \mathbf{C})} \mathbf{M} &\longrightarrow \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{M}^1 \\ &+ \mathbf{M}^{2\dagger} \frac{1}{\omega - E} \mathbf{M}^1 + \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{M}^2 + \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{C} \frac{1}{\omega - E} \mathbf{M}^1 \\ &+ \mathbf{M}^{3\dagger} \frac{1}{\omega - E} \mathbf{M}^1 + \mathbf{M}^{2\dagger} \frac{1}{\omega - E} \mathbf{M}^2 + \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{C} \frac{1}{\omega - E} \mathbf{C} \frac{1}{\omega - E} \mathbf{M}^1 + \dots \end{aligned}$$

→ from here one reads the minimal approximation to \mathbf{C} needed to reproduce the 3rd order diagram. Then the full ladder and ring summation come automatically, for free!

Working eqs. for *ext-ADC(2)*

The matrices for the *extended-ADC(2)* equations are the same as for *ADC(2)*, except for:

$$\mathbf{C}_{(n_1, n_2, k_3), (n_4, n_5, k_6)} = \frac{1}{2} \langle n_1 n_2 | v | n_4 n_5 \rangle \delta_{k_3, k_6} + \langle n_1 k_3 | v^{ph} | n_4 k_6 \rangle \delta_{n_2, n_5} + \langle n_2 k_3 | v^{ph} | n_5 k_6 \rangle \delta_{n_1, n_4}$$

$$\mathbf{D}_{(k_1, k_2, n_3), (k_4, k_5, n_6)} = -\frac{1}{2} \langle k_1 k_2 | v | k_4 k_5 \rangle \delta_{n_3, n_6} - \langle k_1 n_3 | v^{ph} | k_4 n_6 \rangle \delta_{k_2, k_5} - \langle k_2 n_3 | v^{ph} | k_5 n_6 \rangle \delta_{k_1, k_4}$$

where:

$$\langle n_1 n_2 | v | n_4 n_5 \rangle \equiv \mathcal{X}_\gamma^{n_1} \mathcal{X}_\delta^{n_2} v_{\gamma\delta, \mu\nu} (\mathcal{X}_\mu^{n_4} \mathcal{X}_\nu^{n_5})^*$$

$$\langle n_1 k_3 | v^{ph} | n_4 k_6 \rangle \equiv \mathcal{X}_\alpha^{n_1} \mathcal{Y}_\beta^{k_3} v_{\alpha\delta, \beta\gamma} (\mathcal{X}_\gamma^{n_4} \mathcal{Y}_\delta^{k_6})^*$$

$$\langle k_1 k_2 | v | k_4 k_5 \rangle \equiv (\mathcal{Y}_\gamma^{k_1} \mathcal{Y}_\delta^{k_2})^* v_{\gamma\delta, \mu\nu} \mathcal{Y}_\mu^{k_4} \mathcal{Y}_\nu^{k_5}$$


$$\langle k_1 n_3 | v^{ph} | k_4 n_6 \rangle \equiv (\mathcal{Y}_\alpha^{k_1} \mathcal{X}_\beta^{n_3})^* v_{\alpha\delta, \beta\gamma} \mathcal{Y}_\gamma^{k_4} \mathcal{X}_\delta^{n_6}$$

Any repeated indices are implicitly summed

→ The full ladder and ring summations are generated by these choices of **C** and **D**!

Working eqs. For **ADC(3)**


Requiring that ALL 3rd order Goldstone diagrams are included requires to also extending the coupling matrices:



$$\dots + \mathbf{M}^{2\dagger} \frac{1}{\omega - E} \mathbf{M}^1 + \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{C} \frac{1}{\omega - E} \mathbf{M}^1 + \mathbf{M}^{1\dagger} \frac{1}{\omega - E} \mathbf{M}^2 + \dots$$


Working eqs. For **ADC(3)**

Requiring that ALL 3rd order Goldstone diagrams are included requires to also extending the coupling matrices:



$$\begin{aligned} \sqrt{(2)}\mathbf{M}_{(n_1, n_2, k), \alpha} = & \mathcal{X}_\mu^{n_1} \mathcal{X}_\nu^{n_2} \mathcal{Y}_\lambda^k v_{\mu\nu, \alpha\lambda} + \frac{\mathcal{X}_\sigma^{n_1} \mathcal{X}_\rho^{n_2} v_{\sigma\rho, \gamma\delta} \mathcal{Y}_\gamma^{k_7} \mathcal{Y}_\delta^{k_8}}{2(\varepsilon_{k_7}^- + \varepsilon_{k_8}^- - \varepsilon_{n_1}^+ - \varepsilon_{n_2}^+)} (\mathcal{Y}_\mu^{k_7} \mathcal{Y}_\nu^{k_8})^* \mathcal{Y}_\lambda^k v_{\mu\nu, \alpha\lambda} \\ & + \frac{\mathcal{Y}_\sigma^k \mathcal{X}_\rho^{n_2} v_{\rho\delta, \sigma\gamma} \mathcal{Y}_\gamma^{k_5} \mathcal{X}_\delta^{n_6}}{(\varepsilon_k^- - \varepsilon_{n_2}^+ + \varepsilon_{k_5}^- - \varepsilon_{n_6}^+)} \mathcal{X}_\mu^{n_1} (\mathcal{Y}_\nu^{k_5} \mathcal{X}_\lambda^{n_6})^* v_{\mu\nu, \alpha\lambda} - \frac{\mathcal{Y}_\sigma^k \mathcal{X}_\rho^{n_1} v_{\rho\delta, \sigma\gamma} \mathcal{Y}_\gamma^{k_5} \mathcal{X}_\delta^{n_6}}{(\varepsilon_k^- - \varepsilon_{n_1}^+ + \varepsilon_{k_5}^- - \varepsilon_{n_6}^+)} \mathcal{X}_\mu^{n_2} (\mathcal{Y}_\nu^{k_5} \mathcal{X}_\lambda^{n_6})^* v_{\mu\nu, \alpha\lambda} \end{aligned}$$


$$\begin{aligned} \sqrt{(2)}\mathbf{N}_{(k_1, k_2, n), \alpha} = & (\mathcal{Y}_\mu^{k_1} \mathcal{Y}_\nu^{k_2} \mathcal{X}_\lambda^n)^* v_{\mu\nu, \alpha\lambda} + \frac{(\mathcal{Y}_\sigma^{k_1} \mathcal{Y}_\rho^{k_2})^* v_{\sigma\rho, \gamma\delta} (\mathcal{X}_\gamma^{n_7} \mathcal{X}_\delta^{n_8})^*}{2(\varepsilon_{k_1}^- + \varepsilon_{k_2}^- - \varepsilon_{n_7}^+ - \varepsilon_{n_8}^+)} \mathcal{X}_\mu^{n_7} \mathcal{X}_\nu^{n_8} \mathcal{X}_\lambda^n v_{\mu\nu, \alpha\lambda} \\ & + \frac{(\mathcal{Y}_\sigma^{k_2} \mathcal{X}_\rho^n)^* v_{\sigma\delta, \rho\gamma} (\mathcal{X}_\gamma^{n_5} \mathcal{Y}_\delta^{k_6})^*}{(\varepsilon_{k_2}^- - \varepsilon_n^+ + \varepsilon_{k_6}^- - \varepsilon_{n_5}^+)} (\mathcal{Y}_\mu^{k_1})^* \mathcal{X}_\nu^{n_5} \mathcal{Y}_\lambda^{k_6} v_{\mu\nu, \alpha\lambda} - \frac{(\mathcal{Y}_\sigma^{k_1} \mathcal{X}_\rho^n)^* v_{\sigma\delta, \rho\gamma} (\mathcal{X}_\gamma^{n_5} \mathcal{Y}_\delta^{k_6})^*}{(\varepsilon_{k_1}^- - \varepsilon_n^+ + \varepsilon_{k_6}^- - \varepsilon_{n_5}^+)} (\mathcal{Y}_\mu^{k_2})^* \mathcal{X}_\nu^{n_5} \mathcal{Y}_\lambda^{k_6} v_{\mu\nu, \alpha\lambda} \end{aligned}$$



Beyond ADC(3)...

The general strategy is: expand the self-energy in Feynman/Goldstone diagrams up to order n and then compare to the minimal expansion in terms of matrices C , D and M , N .

For ADC(4), also $3p2h/3h2p$ intermediate states appear:





lp/lh-	ADC(2,3)		ADC(4,5)		...
	2p-1h	2h-1p	3p-2h	3h-2p	
$\epsilon + \Sigma(\omega)$	U^I	U^{II}	U^I	U^{II}	...
	$(K+C)^I$		C^I		
		$(K+C)^{II}$		C^{II}	
			$(K+C)^I$		
				$(K+C)^{II}$	





Results for the pairing model

Correlation energy for the 4-level and 4-fermions pairing model





Results for the pairing model

Correlation energy for the 4-level and 4-fermions pairing model



Results for the pairing model



Correlation energy for the 4-level and 4-fermions pairing model



$$\langle \Phi_k; 1h | H_1 | \Phi_{k_1, k_2}^n; 2h1p \rangle = \left\langle \begin{array}{c} \text{---} E_E \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \uparrow \downarrow \\ \uparrow \downarrow \end{array} \middle| \sum_{p \cdot q} P_p^\dagger P_q \middle| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} E_F \text{---} \\ \uparrow \downarrow \\ \uparrow \downarrow \end{array} \right\rangle = 0$$





Results for the pairing model

Correlation energy for the 4-level and 4-fermions pairing model



$$\langle \Phi_k; 1h | H_1 | \Phi_{k_1, k_2}^n; 2h1p \rangle = \left\langle \begin{array}{c} \text{--- } E_E \text{ ---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \middle| \sum_{p \cdot q} P_p^\dagger P_q \middle| \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \\ \text{---} \end{array} E_F \right\rangle = 0$$



Results for the pairing model





By Alexis Mercenne
& Nathan Parzuchowski

Accuracy of ADC(n) - simple atoms/molecules



ADC(1) \equiv HF



ADC(2) \equiv 2nd ord.




ADC(3) \equiv FTDA
FRPA




98-99% of correlation energy is recovered

Accuracy of ADC(n) - simple atoms/molecules


ADC(1) \equiv HF



ADC(2) \equiv 2nd ord.



ADC(3) \equiv FTDA
FRPA



		Hartree-Fock	Second order	FTDA	FRPA	Experiment [63,64]
He	1s	0.918(+14)	0.9012(-2.5)	0.9025(-1.2)	0.9008(-2.9)	0.9037
Be ²⁺	1s	5.6672(+116)	5.6542(-1.4)	5.6554(-0.2)	5.6551(-0.5)	5.6556
Be	2s	0.3093(-34)	0.3187(-23.9)	0.3237(-18.9)	0.3224(-20.2)	0.3426
Ne	1s	4.733(+200)	4.5892(+56)	4.5439(+11)	4.5405(+8)	4.533
	2p	0.852(+57)	0.752(-41)	0.8101(+17)	0.8037(+11)	0.793
Mg ²⁺	2s	1.931(+149)	1.750(-39)	1.8057(+24)	1.7967(+15)	1.782
	2p	3.0068(+56.9)	2.9217(-28.2)	2.9572(+7.3)	2.9537(+3.8)	2.9499
Mg	2s	4.4827	4.3283	4.3632	4.3589	
	3s	0.253(-28)	0.267(-14)	0.272(-9)	0.280(-1)	0.281
Ar	2p	2.282(+162)	2.117(-3)	2.141(+21)	2.137(+17)	2.12
	3p	0.591(+12)	0.563(-16)	0.581(+2)	0.579(\approx 0)	0.579
σ_{rms} [mH]	3s	1.277(+202)	1.111(+36)	1.087(+12)	1.065(-10)	1.075
	3s		1.840	1.578	1.544	
		81.4	29.3	13.7	10.6	

← ionization energies (atoms)

Accuracy of FRPA for atoms




- Diatomic molecules

	FTDA _c	FRPA _c	CCSD(T)	Expt.
N ₂				
E_0	-109.258	-109.272	-109.276	-
r_0	1.104	1.106	1.119	1.098
I	0.565	0.544	0.602 ^a	0.573
BF				
E_0	-124.365	-124.368	-124.380	-
r_0	1.284	1.285	1.295	1.267
I	0.395	0.402	0.406	-
CO				
E_0	-113.037	-113.048	-113.055	-
r_0	1.130	1.123	1.145	1.128
I	0.503	0.494	0.550 ^a	0.515

^a Only up to CCD

Spectral strength of Neon

Momentum distribution:





[CB, Van Neck, AIP Conf.Proc.1120, 104 ('09)]

FIGURE 2. Hole spectral function (right) and momentum distribution (left) of the Ne atom. The dotted, dashed and dot-dashed lines are the contributions coming from the main 2p, 2s and 1s quasihole peaks seen on the right side.

Results for the N-O-F chains

A. Cipollone, CB, P. Navrátil, Phys. Rev. Lett. **111**, 062501 (2013)
and arXiv:1412.3002 [nucl-th] (2014)





→ $d_{3/2}$ raised by genuine 3NF

→ cf. microscopic shell model [Otsuka et al, PRL**105**, 032501 (2010).]

Results for the N-O-F chains

A. Cipollone, CB, P. Navrátil, Phys. Rev. Lett. **111**, 062501 (2013)




→ 3NF crucial for reproducing binding energies and driplines around oxygen

→ $d_{3/2}$ raised by genuine 3NF

→ cf. microscopic shell model [Otsuka et al, PRL**105**, 032501 (2010).]

Results for the N-O-F chains

A. Cipollone, CB, P. Navrátil, Phys. Rev. Lett. **111**, 062501 (2013)
and arXiv:1412.3002 [nucl-th] (2014)




→ 3NF crucial for reproducing binding energies and driplines around oxygen

→ cf. microscopic shell model [Otsuka et al, PRL**105**, 032501 (2010).]

Results for the oxygen chain

A. Cipollone, CB, P. Navrátil, arXiv:1412.3002 [nucl-th] (2014)










→ Single particle spectra slightly to spread and

→ systematic underestimation of radii

Neutron spectral function of Oxygens

A. Cipollone, CB P. Navrátil, *PRC submitted* (2014)





Nuclear matter project with Green's function and coupled cluster

This week we will start looking into how to calculate nuclear matter with the same methods introduced last week. Some short comments on this:

- We will discretize the continuous momentum space by using a box with periodic boundary conditions (PBC). This will be discussed in much detail by Gaute in the next lecture.
- The setup of the basis and the calculation of the reference HF state is the same for all methods (MBPT, CCM, SCGF...) and so next talk will apply to all projects.
- So set up you code with a general basis infrastructure, that will be separate from the solver...
- Some more comments specific to GF:
 - Once the HF is set up, we will need to build bases for pp and hh configurations (as for CC) but also 2p1h and 2h1p. These are all built very similarly.
 - We will use the ADC(2) approach and later move to extended ADC(2), which requires a relatively small extra effort.
 - The self-energy is diagonal in k-space, which is a great simplification: for each value of momentum, we can diagonalize a part of the Dyson equations independently.
 - The approach is very similar to last week pairing model. However, we will have to deal with technical complications (the more sophisticated basis to handle, with more quantum numbers, the dimension of the Dyson sub-matrices, ecc...). All of this will be discussed little by little.